

A NOTE ON RECTIFIABLE SPACES

FUCAI LIN, CHUAN LIU*, AND SHOU LIN

ABSTRACT. In this paper, we firstly discuss the question: Is l_2^∞ homeomorphic to a rectifiable space or a paratopological group? And then, we mainly discuss locally compact rectifiable spaces, and show that a locally compact rectifiable space with the Souslin property is σ -compact, which gives an affirmative answer to A.V. Arhangel'skii and M.M. Choban's question [On remainders of rectifiable spaces, *Topology Appl.*, 157(2010), 789-799]. Next, we show that a rectifiable space X is strongly Fréchet-Urysohn if and only if X is an α_4 -sequential space. Moreover, we discuss the metrizabilities of rectifiable spaces, which gives a partial answer for a question posed in [18]. Finally, we consider the remainders of rectifiable spaces, which improve some results in [2, 3, 6, 19].

1. INTRODUCTION

Recall that a *topological group* G is a group G with a (Hausdorff) topology such that the product maps of $G \times G$ into G is jointly continuous and the inverse map of G onto itself associating x^{-1} with arbitrary $x \in G$ is continuous. A *paratopological group* G is a group G with a topology such that the product maps of $G \times G$ into G is jointly continuous. A topological space G is said to be a *rectifiable space* [11] provided that there are a surjective homeomorphism $\varphi : G \times G \rightarrow G \times G$ and an element $e \in G$ such that $\pi_1 \circ \varphi = \pi_1$ and for every $x \in G$ we have $\varphi(x, x) = (x, e)$, where $\pi_1 : G \times G \rightarrow G$ is the projection to the first coordinate. If G is a rectifiable space, then φ is called a *rectification* on G . It is well known that rectifiable spaces and paratopological groups are all good generalizations of topological groups. In fact, for a topological group with the neutral element e , then it is easy to see that the map $\varphi(x, y) = (x, x^{-1}y)$ is a rectification on G . However, there exists a paratopological group which is not a rectifiable space; Sorgenfrey line ([13, Example 1.2.2]) is such an example. Also, the 7-dimensional sphere S_7 is rectifiable but not a topological group [25, § 3]. Further, it is easy to see that paratopological groups and rectifiable spaces are all homogeneous.

By a remainder of a space X we understand the subspace $bX \setminus X$ of a Hausdorff compactification bX of X .

In section 3, we show that l_2^∞ is homeomorphic to no rectifiable space or paratopological group, where l_2^∞ is the separable Hilbert space, which extends a result of T. Banach in [9]. In section 4, we mainly discuss locally compact rectifiable spaces, and show that a locally compact and separable rectifiable space is σ -compact, which give an affirmative answer for a question of A.V. Arhangel'skii and M.M. Choban's.

2000 *Mathematics Subject Classification.* 54A25; 54B05; 54E20; 54E35.

Key words and phrases. rectifiable spaces; paratopological groups; locally compact; Fréchet-Urysohn; k -networks; sequential spaces; metrizable; compactifications; remainders.

Supported by the NSFC (No. 10971185) and the Educational Department of Fujian Province (No. JA09166) of China.

* corresponding author.

Moreover, we prove that under the set theory assumption a locally compact rectifiable space with the α_4 -properties is metrizable. In section 5, we show that a rectifiable space X is strongly Fréchet-Urysohn if and only if X is an α_4 -sequential space. In section 6, we mainly discuss the metrizability of rectifiable spaces which have a point-countable k -network. In section 7, we mainly consider the question: When does a Tychonoff rectifiable space G have a Hausdorff compactification bG with a remainder belonging to the class of separable and metrizable spaces?

2. PRELIMINARIES

In [22], E. Pentsak studied the topology of the direct limit $X^\infty = \varinjlim X^n$ of the sequence

$$X \subset X \times X \subset X \times X \times X \subset \cdots,$$

where $(X, \star)$ was a “nice” pointed space and X^n was identified with the subspace $X^n \times \{\star\}$ of X^{n+1} .

A space X is called an S_2 -space (*Arens’ space*) if $X = \{\infty\} \cup \{x_n : n \in \mathbb{N}\} \cup \{x_n(m) : m, n \in \mathbb{N}\}$ and the topology is defined as follows: Each $x_n(m)$ is isolated; a basic neighborhood of x_n is $\{x_n\} \cup \{x_n(m) : m > k\}$ for some $k \in \mathbb{N}$; a basic neighborhood of ∞ is $\{\infty\} \cup (\bigcup \{V_n : n > k\})$ for some $k \in \mathbb{N}$, where V_n is a neighborhood of x_n .

T. Banach define the space K [9].

Let

$$K = \{(0, 0)\} \cup \{(\frac{1}{n}, \frac{1}{nm}) : n, m \in \mathbb{N}\} \subset \mathbb{R}^2.$$

The space K is non-locally compact and metrizable. Also, the space K is a minimal space with these properties in the sense that each metrizable non-locally compact space contains a closed copy of K . For convenience, put $x_0 = (0, 0)$ and $x_{n,m} = (\frac{1}{n}, \frac{1}{nm})$ for any $n, m \in \mathbb{N}$.

The space X is called S_ω if X is obtained by identifying all the limit points of ω many convergent sequences.

If A is a subset of a space X , then $[A]^{Seq}$ denotes the *sequential closure* of A , i.e. the set of limits of convergent sequences in A . Clearly, we have $A \subset [A]^{Seq}$. By induction on $\alpha \in \omega_1 + 1$, we can define $[A]_\alpha$ as follows: $[A]_0 = A$, $[A]_{\alpha+1} = [[A]_\alpha]^{Seq}$ and $[A]_\alpha = \bigcup \{[A]_\beta : \beta < \alpha\}$ for a limit order α . One can easily verify that $[A]_{\omega_1+1} = [A]_{\omega_1}$, and that a space X is sequential iff $\overline{A} = [A]_{\omega_1}$ for every $A \subset X$. For a sequential space X we define $so(X)$, the *sequential order* of X , by $so(X) = \min\{\alpha \in \omega_1 + 1 : \overline{A} = [A]_\alpha \text{ for each } A \subset X\}$.

Definition 2.1. A space X is said to be *Fréchet-Urysohn* if, for each $x \in \overline{A} \subset X$, there exists a sequence $\{x_n\}$ such that $\{x_n\}$ converges to x and $\{x_n : n \in \mathbb{N}\} \subset A$. A space X is said to be *strongly Fréchet-Urysohn* if the following condition is satisfied

(SFU) For every $x \in X$ and each sequence $\eta = \{A_n : n \in \mathbb{N}\}$ of subsets of X such that $x \in \bigcap_{n \in \mathbb{N}} \overline{A_n}$, there is a sequence $\zeta = \{a_n : n \in \mathbb{N}\}$ in X converging to x and intersecting infinitely many members of η .

Obviously, a strongly Fréchet-Urysohn space is Fréchet-Urysohn. However, the space S_ω is Fréchet-Urysohn and non-strongly Fréchet-Urysohn.

Let X be a space. For $P \subset X$, the set P is a *sequential neighborhood* of x in X if every sequence converging to x is eventually in P .

Definition 2.2. Let $\mathcal{P} = \bigcup_{x \in X} \mathcal{P}_x$ be a cover of a space X such that for each $x \in X$, (a) if $U, V \in \mathcal{P}_x$, then $W \subset U \cap V$ for some $W \in \mathcal{P}_x$; (b) the family $\mathcal{P}_x$ is a network of x in X , i.e., $x \in \bigcap \mathcal{P}_x$, and if $x \in U$ with U open in X , then $P \subset U$ for some $P \in \mathcal{P}_x$.

The family $\mathcal{P}$ is called a *weak base* for X [1] if, for every $G \subset X$, the set G must be open in X whenever for each $x \in G$ there exists $P \in \mathcal{P}_x$ such that $P \subset G$. The space X is *weakly first-countable* if the family $\mathcal{P}$ is a weak base for X such that each $\mathcal{P}_x$ is countable.

The following theorem for the first time there was announced in [11], and the readers can see the proof in [12, 15, 24].

Theorem 2.3. [11] *A topological space G is rectifiable if and only if there exists $e \in G$ and two continuous maps $p : G^2 \rightarrow G$, $q : G^2 \rightarrow G$ such that for any $x \in G, y \in G$ the next identities hold:*

$$p(x, q(x, y)) = q(x, p(x, y)) = y, q(x, x) = e.$$

In fact, we can assume that $p = \pi_2 \circ \varphi^{-1}$ and $q = \pi_2 \circ \varphi$ in Theorem 2.3. Fixed a point $x \in G$, then $f_x, g_x : G \rightarrow G$ defined with $f_x(y) = p(x, y)$ and $g_x(y) = q(x, y)$, for each $y \in G$, are homeomorphism, respectively. We denote f_x, g_x with $p(x, G), q(x, G)$, respectively.

Let G be a rectifiable space, and let p be the multiplication on G . Further, we sometime write $x \cdot y$ instead of $p(x, y)$ and $A \cdot B$ instead of $p(A, B)$ for any $A, B \subset G$. Therefore, $q(x, y)$ is an element such that $x \cdot q(x, y) = y$; since $x \cdot e = x \cdot q(x, x) = x$ and $x \cdot q(x, e) = e$, it follows that e is a right neutral element for G and $q(x, e)$ is a right inverse for x . Hence a rectifiable space G is a topological algebraic system with operation p, q , 0-ary operation e and identities as above. It is easy to see that this algebraic system need not to satisfy the associative law about the multiplication operation p . Clearly, every topological loop is rectifiable.

All spaces are T_1 and regular unless stated otherwise. The notation $\mathbb{N}$ denotes the set of all positive natural numbers. The letter e denotes the neutral element of a group and the right neutral element of a rectifiable space, respectively. Readers may refer to [4, 13, 14] for notations and terminology not explicitly given here.

3. l_2^∞ IS HOMEOMORPHIC TO NO RECTIFIABLE SPACE OR PARATOPOLOGICAL GROUPS

In this section, by a modification of the proof of Theorem 1 in [9], we show that l_2^∞ is homeomorphic to no rectifiable space or paratopological group.

We call a subset A of a rectifiable space (resp. paratopological group) G *multiplicative* if for any $a, b \in A$ we have $p(a, b) = a \cdot b \in A$ (resp. $ab \in A$).

We denote by $\text{conv}(K) = \{(0, 0)\} \cup \{(x, y) : 0 < y \leq x \leq 1\}$, where $\text{conv}(K)$ is the convex hull of K in $\mathbb{R}^2$.

In this section, we may assume that $S_\omega = \{y_0\} \cup \{y_{n,m} = (n, \frac{1}{m}) : n, m \in \mathbb{N}\}$, where, for each $n \in \mathbb{N}$, the sequence $\{y_{n,m}\} \rightarrow y_0$ as $m \rightarrow \infty$. For each $k \in \mathbb{N}$, let $V_k = \{y_0\} \cup \{y_{n,m} : n \leq k, m \in \mathbb{N}\}$. It is easy to see that S_ω has the direct limit topology with respect to the sequence $V_1, V_2, \dots$.

Theorem 3.1. *Let X be a normal k -space. If X contains closed copies of S_ω and K , then it is homeomorphic to no closed multiplicative subset of a rectifiable space G such that y_0 is the right neutral element of G .*

Proof. Suppose not, let X be a closed multiplicative subset of a rectifiable space G . Now, we define a map $f : K \times S_\omega \rightarrow X$ with $f(x, y) = p(x, y)$ for each $(x, y) \in K \times S_\omega$. Then the following (1) and (2) hold:

- (1) the map $(\pi_1, f) : K \times S_\omega \rightarrow K \times X$ is a closed embedding, where $(\pi_1, f)(x, y) = (x, f(x, y))$ for each $(x, y) \in K \times S_\omega$;
- (2) the map $g : K \rightarrow X$ defined by $g(x) = p(x, y_0) = x$ (that is, g is the identity map), for each $x \in K$, is a closed embedding.

Indeed, the statement (2) is obvious. Moreover, it is easy to see that the map $(\pi_1, f) : K \times S_\omega \rightarrow K \times X$ is injective continuous. We only show that the map (π_1, f) is relatively open. For each open subset $U \times V$ of $K \times S_\omega$, we have $(\pi_1, f)(U \times V) = \cup\{\{x\} \times (x \cdot V) : x \in U\}$, where U and V are open in K and S_ω respectively. Since V is open in S_ω , there exists an open subset W of X such that $W \cap S_\omega = V$. Therefore, we have

$$(\pi_1, f)(U \times V) = \cup\{\{x\} \times p(x, V) : x \in U\} = (U \times (U \cdot W)) \cap (\pi_1, f)(K \times S_\omega).$$

Since $p(U, W) = \cup\{p(x, W) : x \in U\}$ is open in X , the set $U \times p(U, W)$ is open in $K \times X$.

By the normality of X , let $h : X \rightarrow \text{conv}(K)$ be a continuous extension of the map $g^{-1} : g(K) \rightarrow K$.

For each $n, m \in \mathbb{N}$, let $\delta_{n,m} = \frac{1}{2nm(m+1)}$, and put

$$W_{n,m} = \text{conv}(K) \cap ((\frac{1}{n} - \delta_{n,m}, \frac{1}{n} + \delta_{n,m}) \times (\frac{1}{nm} - \delta_{n,m}, \frac{1}{nm} + \delta_{n,m})).$$

Obviously, the collection $\{W_{n,m} : n, m \in \mathbb{N}\}$ consists of pairwise disjoint neighborhoods of the points $x_{n,m}$ in $\text{conv}(K)$. Since $y_{n,m} \rightarrow y_0$ as $m \rightarrow \infty$ and $h \circ f(x_{n,m}, y_0) = h(p(x_{n,m}, y_0)) = x_{n,m} = (\frac{1}{n}, \frac{1}{nm})$, for any $n, m \in \mathbb{N}$, there exists a $k(n, m) \in \mathbb{N}$ such that $h \circ f(x_{n,m}, y_{n,k(n,m)}) \in W_{n,m}$. Without loss of generality, we may assume that $k(n, m+1) > k(n, m)$ for any $n, m \in \mathbb{N}$. Put

$$Z = \{p(x_{n,m}, y_{n,k(n,m)}) : n, m \in \mathbb{N}\}.$$

For each $n, m \in \mathbb{N}$, it follows from $h \circ f(x_0, y_0) \notin W_{n,m}$ that $f(x_0, y_0) \notin Z$.

Claim: Z is closed in X .

Since X is a k -space, it suffices to prove that for each compact subset F of X the intersection $F \cap Z$ is closed in F . Let

$$F_1 = \{x_0\} \cup \{x_{n,m} : h(F) \cap W_{n,m} \neq \emptyset, n, m \in \mathbb{N}\} \text{ and } F_2 = \pi_2((\pi_1, f)^{-1}(F_1 \times F)).$$

Since $h(F) \subset \text{conv}(K)$ is compact, the set F_1 is compact. It follows from (1) that

$$(\pi_1, f)^{-1}(F \times F_1) \subset K \times S_\omega$$

is compact, and hence F_2 is also compact. Because $S_\omega = \varinjlim V_n$, there exists an $n_0 \in \mathbb{N}$ such that $F_2 \subset V_{n_0}$. Since $F \cap Z \subset p(F_1, F_2)$, we have $F \cap Z \subset \{p(x_{n,m}, y_{n,k(n,m)}) : n \leq n_0, y_{n,m} \in F_1\}$. By the compactness of F_1 , it is easy to see that $\{p(x_{n,m}, y_{n,k(n,m)}) : n \leq n_0, x_{n,m} \in F_1\}$ is finite. Therefore, the set $F \cap Z$ is closed in F .

Since $p(x_0, y_0) = x_0 \notin Z$ and p is continuous, it follows from the Claim that there exist open neighborhoods $V(x_0) \subset K$ and $U(y_0) \subset S_\omega$ of x_0 and y_0 respectively such that $p(V(x_0), U(y_0)) \cap Z = \emptyset$. For every $m \in \mathbb{N}$, we can fix an $n \in \mathbb{N}$ such that $x_{n,m} \in V(x_0)$. Since $\{y_{n,m}\}_{m=1}^\infty$ converges to y_0 and $\{k(n, m)\}_{m=1}^\infty$ is increasing, there is an $m \in \mathbb{N}$ such that $y_{n,k(n,m)} \in U(y_0)$. Then $p(x_{n,m}, y_{n,k(n,m)}) \in p(V(x_0), U(y_0)) \cap Z$, which is a contradiction. $\square$

Theorem 3.2. *Let X be a normal k -space. If X contains closed copies of S_ω and K , then it is homeomorphic to no closed multiplicative subset of a paratopological group.*

Proof. Suppose not, let X be a closed multiplicative subset of a rectifiable space G . Now, we define a map $f : K \times S_\omega \rightarrow X$ with $f(x, y) = xy$ for each $(x, y) \in K \times S_\omega$. Obviously, we can obtain the following (1) and (2):

(1) the map $(\pi_1, f) : K \times S_\omega \rightarrow K \times X$ is a closed embedding, where $(\pi_1, f)(x, y) = (x, xy)$ for each $(x, y) \in K \times S_\omega$;

(2) the map $g : K \rightarrow X$ defined by $g(x) = xy_0$, for each $x \in K$, is a closed embedding.

By the normality of X , let $h : X \rightarrow \text{conv}(K)$ be a continuous extension of the map $g^{-1} : g(K) \rightarrow K$.

By the proof of Theorem 3.1, we can define the neighborhoods $W_{n,m}$ of the points $x_{n,m}$ in $\text{conv}(K)$ and the closed set Z with $x_0y_0 \notin Z$.

Since $f(x_0, y_0) = x_0y_0 \notin Z$, G is joint continuous and Z is closed, it follows that there exist open neighborhoods $V(x_0) \subset K$ and $U(y_0) \subset S_\omega$ of x_0 and y_0 respectively such that $(V(x_0) \times U(y_0)) \cap Z = \emptyset$. For every $m \in \mathbb{N}$, we can fix an $n \in \mathbb{N}$ such that $x_{n,m} \in V(x_0)$. Since $\{y_{n,m}\}_{m=1}^\infty$ converges to y_0 and $\{k(n, m)\}_{m=1}^\infty$ is increasing, there is an $m \in \mathbb{N}$ such that $y_{n,k(n,m)} \in U(y_0)$. Then $x_{n,m}y_{n,k(n,m)} \in (V(x_0) \times U(y_0)) \cap Z$, which is a contradiction. $\square$

It is well known that a space X contains a closed copy of S_ω , provided X can be written as a direct limit of a sequence

$$X_1 \subset X_2 \subset \cdots,$$

where each X_n is a closed metrizable subset of X , nowhere dense in X_{n+1} . In particular, the space l_2^∞ contains a topological closed copy of S_ω . Moreover, the space l_2^∞ is a normal k -space and contains a topological closed copy of K . Therefore, by the topological homogeneity of l_2^∞ and Theorems 3.1 and 3.2, we have the following theorem.

Theorem 3.3. *l_2^∞ is homeomorphic to no rectifiable space or a paratopological group.*

Corollary 3.4. *l_2^∞ is homeomorphic to no topological loop.*

Corollary 3.5. [9] *l_2^∞ is homeomorphic to no topological group.*

4. LOCALLY COMPACT RECTIFIABLE SPACES

In [6], A.V. Arhangel'skiĭ and M.M. Choban posed the following question:

Question 4.1. [6, Problem 5.10] *Is every rectifiable p -space with a countable Souslin number Lindelöf? What if we assume the space to be separable? Separable and locally compact?*

Now, we give an affirmative answer for Questions 4.1 of the case of separable and locally compact rectifiable spaces.

Lemma 4.2. *Let G be a rectifiable space. If Y is a dense subset of G and U is an open neighborhood of the right neutral element e of G , then $G = Y \cdot U$.*

Proof. Fix an arbitrary $z \in G$. Since $q(z, z) = e \in U$, there exists an open neighborhood V of e such that $q(z \cdot V, z) \subset U$. Put $W = z \cdot V$. Then W is an open neighborhood of z in G . Since Y is a dense subset of G , we have $W \cap Y \neq \emptyset$. Take a point $y \in W \cap Y = z \cdot V \cap Y$. Then $y = z \cdot v$ for some $v \in V$.

$$z = p(z \cdot v, q(z \cdot v, z)) = p(y, q(z \cdot v, z)) \in p(y, q(z \cdot V, z)) \subset p(y, U) = y \cdot U \subset Y \cdot U.$$

By the choice of arbitrary point of z , we have $G = Y \cdot U$. $\square$

It follows from Lemma 4.2, we have the following results, which give an answer for Question 4.1.

Theorem 4.3. *If G is a locally σ -compact¹ rectifiable space with the Souslin property, then G is σ -compact.*

Proof. Let G be a locally σ -compact rectifiable space with the Souslin property. For each $\alpha \in \Gamma$, let $\mathcal{A}_\alpha$ be the family consisting of disjoint open subsets of G such that each element of $\mathcal{A}_\alpha$ is covered by countably many compact subsets (since G is locally σ -compact). $\{\mathcal{A}_\alpha, \alpha \in \Gamma\}$ is a set with partial order by inclusion. It is easy to see that every chain of $\{\mathcal{A}_\alpha, \alpha \in \Gamma\}$ has an upper bound, by Zorn's Lemma, there is a maximal element $\mathcal{A} \in \{\mathcal{A}_\alpha, \alpha \in \Gamma\}$. Since G has Souslin property, we have $|\mathcal{A}| \leq \omega$, and hence we write $\mathcal{A} = \{A_i\}$, $A_i \subset \cup\{K_{i,j}\}$, where each $K_{i,j}$ is a compact subset of G . By maximality of $\mathcal{A}$, $\cup\{K_{i,j}\}$ is a dense subset of G . Let U be an open neighborhood of e , which is covered by countably many compact subsets $\{H_l\}$. By Lemma 4.2, $G = (\cup\{K_{i,j}\}) \cdot (\cup\{H_l\}) = \cup(K_{i,j} \cdot H_l)$, each $K_{i,j} \cdot H_l$ is compact, hence G is σ -compact. $\square$

Corollary 4.4. *If G is a locally compact and separable rectifiable space, then G is σ -compact (and, hence, Lindelöf).*

Corollary 4.5. *If G is a locally Lindelöf and separable rectifiable space, then G is Lindelöf.*

Let A be a subspace of a rectifiable space G . Then A is called a *rectifiable subspace* of G if we have $p(A, A) \subset A$ and $q(A, A) \subset A$.

Proposition 4.6. *Let G be a rectifiable space. If H is a rectifiable subspace of G , then $\overline{H}$ is also a rectifiable subspace of G .*

Proof. Take two points $x, y \in \overline{H}$. Then we shall show that $p(x, y) \in \overline{H}$ and $q(x, y) \in \overline{H}$.

Since $x, y \in \overline{H}$, there exist two nets $\{x_\alpha\}, \{y_\beta\}$ in H such that $x_\alpha \rightarrow x, y_\beta \rightarrow y$. Since p is continuous, $p(x, y)$ is a cluster point of $\{p(x_\alpha, y_\beta) \subset H$. Hence $p(x, y) \in \overline{H}$.

Similarly, we can show that $q(x, y) \in \overline{H}$. $\square$

Lemma 4.7. *Let G be a rectifiable space. If V is an open rectifiable subspace of G , then V is closed in G .*

Proof. Suppose that V is non-closed in G . Then $\overline{V} \setminus V \neq \emptyset$. Take a point $x \in \overline{V} \setminus V$. Since $q(x, x) = e \in V$ and the continuity of q , there exists an open neighborhood W of e such that $q(x \cdot W, x) \subset V$. Put $U = x \cdot W$. Then U is an open neighborhood

¹A space X is *locally σ -compact* if, for each point x of X , there exists an open neighborhood U_x of x such that U_x can be cover by a countably many compact subsets of X .

of x , and hence $U \cap V \neq \emptyset$ since $x \in \overline{V}$. Therefore, there exist $a \in W$ and $b \in V$ such that $x \cdot a = b$. Then we have

$$x = p(x \cdot a, q(x \cdot a, x)) = p(b, q(x \cdot a, x)) \subset p(V, V) = V,$$

where $p(V, V) = V$ since V is a rectifiable subspace of G . However, the point $x \notin V$, which is a contradiction. $\square$

Theorem 4.8. *If H is a locally compact rectifiable subspace of a rectifiable space G , then H is closed in G .*

Proof. Let $K = \overline{H}$. Then K is a rectifiable subspace of G by Proposition 4.6. Since H is a dense locally compact subspace of K , it follows from [13, Theorem 3.3.9] that H is open in K . By Lemma 4.7, the set H is closed, and hence $K = H$. $\square$

The following Lemma maybe was proved somewhere.

Lemma 4.9. *Let F be a compact subset of a space X and have a countable base $\{U_n\}$ with $\overline{U_{n+1}} \subset U_n$ in X , and let $H = \bigcap_n V_n$ ($V_{n+1} \subset V_n$ and each V_n is open in F) is a compact G_δ -set of F . For $n \in \mathbb{N}$, let W_n be an open set in X such that $V_n = W_n \cap F$, $W_n \subset U_n$, $\overline{W_{n+1}} \subset W_n$, then $\{W_n\}$ is a countable base at H in X .*

Proof. $H = \bigcap_n W_n = \bigcap_n \overline{W_n}$. Suppose that $\{W_n\}$ is not a countable base at H , then there is an open subset U of X such that $H \subset U$ and $W_n \setminus U \neq \emptyset$. By induction, choose $x_n \in W_n \setminus U$ with $x_i \neq x_j$ if $i \neq j$. Since $x_n \in U_n$ for each $n \in \mathbb{N}$, then $\{x_n\}$ has a cluster point x . In fact, if $\{x_n\} \cap F$ is infinite, then $\{x_n\}$ has a cluster point in F since F is compact; if $\{x_n\} \cap F$ is finite, without loss generality, we assume $\{x_n\} \cap F = \emptyset$. Since $F \subset X \setminus \{x_n\}$ which is open in X , there is $n_0 \in \mathbb{N}$ such that $F \subset U_{n_0} \subset X \setminus \{x_n\}$ for $n > n_0$. This is a contradiction since $x_n \in U_n$. Therefore, we have $x \in \overline{W_n}$ for each n , then $x \in H \subset U$, and hence U contains infinitely many x'_n s, which is a contradiction. $\square$

Next, we shall show that, for each locally compact rectifiable space, there exists a compact rectifiable subspace with a countable character.

Lemma 4.10. *Let G be a rectifiable space and F be a compact subset of G containing e and having a countable base $\{U_n : n \in \mathbb{N}\}$ in G . Assume that a sequence $\zeta = \{V_n : n \in \mathbb{N}\}$ of open neighborhoods of e in G such that $\overline{V_{n+1}} \cdot \overline{V_{n+1}} \subset V_n \cap U_n$ and $q(V_{n+1}, V_{n+1}) \subset V_n$. Then $H = \bigcap_{n \in \mathbb{N}} V_n$ is a compact rectifiable subspace of G , $H \subset F$ and ζ is a base for G at H .*

Proof. Obviously, we have $\overline{V_{n+1}} \subset V_n$ for each $n \in \mathbb{N}$. We first claim that H is a compact rectifiable subspace of G .

Indeed, it is easy to see that $H = \bigcap_{n \in \mathbb{N}} V_n = \bigcap_{n \in \mathbb{N}} \overline{V_n}$, and hence H is closed in G . For each $x, y \in H$, we have $x, y \in V_n$ for each $n \in \mathbb{N}$. Then, for each $n \in \mathbb{N}$, we have $x \cdot y \in V_n$ since $\overline{V_{n+1}} \cdot \overline{V_{n+1}} \subset V_n$. Therefore, $x \cdot y \in H$. Since $q(V_{n+1}, V_{n+1}) \subset V_n$, we have $q(x, y) \in H$. Therefore, H is a rectifiable closed subspace. Obviously, $H \subset \bigcap_{n \in \mathbb{N}} U_n = F$. Thus H is compact. By Lemma 4.9, ζ is a base for G at H . $\square$

Proposition 4.11. *Let G be a rectifiable space with point-countable type. If O is an open neighborhood of e , then there exists a compact rectifiable subspace H of countable character in G satisfying $H \subset O$.*

Proof. Since G is of point-countable type, there exists a compact subset of G having a countable base in G . By the homogeneity of G , we may assume that $e \in F$. Let $\{U_n : n \in \mathbb{N}\}$ be a countable base for G at F . We define by induction a sequence $\{V_n : n \in \mathbb{N}\}$ of open neighborhoods of e in G satisfying the following conditions:

- (1) $V_1 \subset O$;
- (2) $\overline{V_{n+1}} \cdot \overline{V_{n+1}} \subset V_n \cap U_n$ for each $n \in \mathbb{N}$;
- (3) $q(V_{n+1}, V_{n+1}) \subset V_n$ for each $n \in \mathbb{N}$.

Put $H = \bigcap_{n \in \mathbb{N}} V_n$. Then $H \subset O$ by (1). It follows from Lemma 4.10 that H is a compact rectifiable subspace of G and that $\{V_n : n \in \mathbb{N}\}$ is a base of G at H . $\square$

Since each locally compact space is of point-countable type, we have the following corollary.

Corollary 4.12. *Let G be a locally compact rectifiable space. If O is an open neighborhood of e , then there exists a compact rectifiable subspace H of countable character in G satisfying $H \subset O$.*

Definition 4.13. Let X be a topological space. For $i = 1, 4$ we say that X is an α_i -space if for each countable family $\{S_n : n \in \mathbb{N}\}$ of sequences converging to some point $x \in X$ there is a sequence S converging to x such that:

- (α_1) $S_n \setminus S$ is finite for all $n \in \mathbb{N}$;
- (α_4) $S_n \cap S \neq \emptyset$ for infinitely many $n \in \mathbb{N}$.

Obviously, we have $\alpha_1 \Rightarrow \alpha_4$.

Let ω^ω denote the family of all functions from $\mathbb{N}$ into $\mathbb{N}$. For $f, g \in \omega^\omega$ we write $f <^* g$ if $f(n) < g(n)$ for all but finitely many $n \in \mathbb{N}$. A family $\mathcal{F}$ is *bounded* if there is a $g \in \omega^\omega$ such that $f <^* g$ for all $f \in \mathcal{F}$, and is *unbounded* otherwise. We denote by $\mathfrak{b}$ the smallest cardinality of an unbounded family in ω^ω . It is easy to see that $\omega < \mathfrak{b} \leq \mathfrak{c}$, where $\mathfrak{c}$ denotes the cardinality of the continuum.

Lemma 4.14. [21] *For $i \in \{1, 4\}$, D^τ is an α_i -space if and only if $\tau < \mathfrak{b}$, where D is the discrete two-points space $\{0, 1\}$.*

Theorem 4.15. *The following conditions are equivalent:*

- (1) *Every compact rectifiable space with the α_1 -property is metrizable;*
- (2) *Every locally compact rectifiable space with the α_4 -property is metrizable;*
- (3) $\mathfrak{b} = \omega_1$.

Proof. The implication (2) $\Rightarrow$ (1) is trivial.

(1) $\Rightarrow$ (3). Since D^{ω_1} is a nonmetrizable compact group, so it cannot be an α_1 -space by (1). It follows from Lemma 4.14 that $\mathfrak{b} \leq \omega_1$. Since $\mathfrak{b} > \omega$, it follows that $\mathfrak{b} = \omega_1$.

(3) $\Rightarrow$ (2). Suppose that $\mathfrak{b} = \omega_1$, and that G is a locally compact α_4 -rectifiable space. Next, we shall prove that G is metrizable. By Proposition 4.11, there exists a compact rectifiable subspace F of G which has a countable character at F in G . We claim that F is metrizable. If not, as proved V.V. Uspenskij in [24, 25], compact rectifiable spaces are dyadic, and hence the space F contains a subspace homeomorphic to D^{ω_1} . Since a subspace of an α_4 -space is an α_4 -space, the subspace D^{ω_1} is an α_4 -space. Then, it follows from Lemma 4.14 that $\omega_1 < \mathfrak{b}$, which is a contradiction. Therefore, the space F is metrizable.

Let $\{U_n : n \in \mathbb{N}\}$ be a countable base of G at F , where $\overline{U_{n+1}} \subset U_n$ for each $n \in \mathbb{N}$. Let $\{V_n : n \in \mathbb{N}\}$ be a countable neighborhoods base at the point e in F ,

where the closure $\text{cl}_F V_{n+1} \subset V_n$ for each $n \in \mathbb{N}$. For each $n \in \mathbb{N}$, there exists an open subset W_n of G such that $V_n = W_n \cap F$, $W_n \subset \overline{U_n}$ and $\overline{W_{n+1}} \subset W_n$. Put $\gamma = \{W_n : n \in \mathbb{N}\}$. By Lemma 4.9, the family γ is a neighborhood base in G at point e . hence the space G is first-countable, and therefore, it is metrizable. $\square$

Corollary 4.16. [21] *The following conditions are equivalent:*

- (1) *Every compact topological group with the α_1 -property is metrizable;*
- (2) *Every locally compact topological group with the α_4 -property is metrizable;*
- (3) $\flat = \omega_1$.

Question 4.17. *Let G be a locally compact rectifiable α_4 -space. Is the space G an α_1 -space in ZFC?*

5. α_4 -RECTIFIABLE SPACES

In this section, we first give a new proof of the properties of Fréchet-Urysohn and strongly Fréchet-Urysohn are coincide in rectifiable spaces, which was proved in [18].

First, we recall a concept.

(AS) For any family $\{a_{m,n} : (m,n) \in \mathbb{N} \times \mathbb{N}\} \subset X$ with $\lim_n a_{m,n} = a \in X$ for each $m \in \mathbb{N}$, it is possible to choose two strictly increasing sequences $\{i_l\}_{l \in \mathbb{N}} \subset \mathbb{N}$ and $\{j_l\}_{l \in \mathbb{N}} \subset \mathbb{N}$ such that $\lim_l a_{i_l, j_l} = a$. Obviously, a space with AS-property is an α_4 -space.

It is well known that a topological space X is a strongly Fréchet-Urysohn space if and only if it is Fréchet-Urysohn and has the double sequence property (α_4). Therefore, it is sufficient to show that a Fréchet-Urysohn rectifiable space has the double sequence property (α_4). Indeed, we have the following result.

Lemma 5.1. *A Fréchet-Urysohn Hausdorff rectifiable space G satisfies AS and hence α_4 as well.*

Proof. Assume that G is a non-discrete space. Let $\{a_{m,n} : (m,n) \in \mathbb{N} \times \mathbb{N}\} \subset X$ with $\lim_n a_{m,n} = e$ for each $m \in \mathbb{N}$. Since G is a Fréchet-Urysohn non-discrete space, there exists a sequence $\{s_m\}_{m \in \mathbb{N}} \subset G$ with $\lim_m s_m = e$ such that $s_m \neq e$ for each $m \in \mathbb{N}$.

Put $z_{m,k} = q(s_m, a_{m,k+m})$ if $q(s_m, a_{m,k+m}) \neq e$, and $z_{m,k} = s_m$ if $q(s_m, a_{m,k+m}) = e$. Let $M = \{z_{m,k} : (m,k) \in \mathbb{N} \times \mathbb{N}\}$. Obviously, we have $e \notin M$ since $s_m \neq e$ for each $m \in \mathbb{N}$. However, we have $e \in \overline{M}$. Indeed, if $M \cap \{s_m : m \in \mathbb{N}\}$ is infinite, then it is easy to see that $e \in \overline{M}$. Therefore, suppose that $M \cap \{s_m : m \in \mathbb{N}\}$ is finite. Then there is an open neighborhood U of e such that $U \cap M \cap \{s_m : m \in \mathbb{N}\} = \emptyset$. Let V be any open neighborhood of e with $V \subset U$. Hence there is an open neighborhood W of e such that $q(W, W) \subset V$. It follows from $\lim_m s_m = e$ that there exists an $m \in \mathbb{N}$ such that $s_m \in W$. Since $\lim_n a_{m,n} = e$, there exists a $k \in \mathbb{N}$ such that $a_{m,k+m} \in W$. Therefore, we have $q(s_m, a_{m,k+m}) = z_{m,k} \in q(W, W) \subset V \subset U$.

Since $e \in \overline{M}$ and G is Fréchet-Urysohn, we can find a sequence $\{(m_l, k_l)\}_{l \in \mathbb{N}}$ in G such that $\lim_l z_{m_l, k_l} = e$.

Case 1: The sequence $\{k_l\}_{l \in \mathbb{N}}$ is bounded.

Without loss of generality, we may assume that $k_l = r$ for each $l \in \mathbb{N}$ for some $r \in \mathbb{N}$. Since $\lim_l z_{m_l, k_l} = \lim_l z_{m_l, r} = e$ and $z_{m_l, r} \neq e$ for each $l \in \mathbb{N}$, we have

$\lim_l m_l = \infty$. Without loss of generality, suppose that $m_l < m_{l+1}$ for each $l \in \mathbb{N}$. Let $N_1 = \{l \in \mathbb{N} : z_{m_l, r} = s_{m_l}\}$.

Subcase 1.1: The set N_1 is infinite.

We denote N_1 by $\{p_i : i \in \mathbb{N}\}$, where $p_i < p_{i+1}$ for each $i \in \mathbb{N}$. Then it is easy to see that $q(s_{m_{p_l}}, a_{m_{p_l}, r+m_{p_l}}) = e$ for each $l \in \mathbb{N}$. Since $\lim_l s_{m_{p_l}} = e$, we have

$$a_{m_{p_l}, r+m_{p_l}} = p(s_{m_{p_l}}, q(s_{m_{p_l}}, a_{m_{p_l}, r+m_{p_l}})) = p(s_{m_{p_l}}, e) = s_{m_{p_l}} \rightarrow e \text{ as } l \rightarrow \infty.$$

Therefore, we can set $i_l = m_{p_l}$ and $j_l = r + m_{p_l}$ for each $l \in \mathbb{N}$. Then we get the strictly increasing sequences $\{i_l\}_{l \in \mathbb{N}}$ and $\{j_l\}_{l \in \mathbb{N}}$ such that $\lim_l a_{i_l, j_l} = e$.

Subcase 1.2: The set N_1 is finite.

Let $N_2 = \{l \in \mathbb{N} : z_{m_l, r} \neq s_{m_l}\}$. Then N_2 is infinite. We may denote N_2 by $\{q_i : i \in \mathbb{N}\}$, where $q_i < q_{i+1}$ for each $i \in \mathbb{N}$. It follows that

$$z_{m_{q_l}, k_{q_l}} = q(s_{m_{q_l}}, a_{m_{q_l}, r+m_{q_l}}) \text{ for each } l \in \mathbb{N}.$$

Since $\lim_l z_{m_{q_l}, k_{q_l}} = e$ and $\lim_l s_{q_l} = e$, we have

$$a_{m_{q_l}, r+m_{q_l}} = p(s_{q_l}, q(s_{m_{q_l}}, a_{m_{q_l}, r+m_{q_l}})) = p(s_{q_l}, z_{m_{q_l}, k_{q_l}}) \rightarrow p(e, e) = e \text{ as } l \rightarrow \infty.$$

Therefore, we can set $i_l = m_{q_l}$ and $j_l = r + m_{q_l}$ for each $l \in \mathbb{N}$. Then we get the strictly increasing sequences $\{i_l\}_{l \in \mathbb{N}}$ and $\{j_l\}_{l \in \mathbb{N}}$ such that $\lim_l a_{i_l, j_l} = e$.

Case 2: The sequence $\{k_l\}_{l \in \mathbb{N}}$ is unbounded.

Without loss of generality, we may assume that $\{k_l\}_{l \in \mathbb{N}}$ is a strictly increasing sequence.

Claim: $\lim_l m_l = \infty$.

If not, we may assume that, for each $l \in \mathbb{N}$, $m_l = t$ for some $t \in \mathbb{N}$. Since $\{k_l\}_{l \in \mathbb{N}}$ is strictly increasing, we have $\lim_l a_{t, t+k_l} = e$. It follows from $\lim_l z_{t, k_l} = e$ that

$$a_{t, t+k_l} = a_{m_l, m_l+k_l} = p(s_{m_l}, z_{m_l+k_l}) = p(s_t, z_{t, k_l}) \rightarrow p(s_t, e) = s_t \text{ as } l \rightarrow \infty.$$

However, $a_{t, t+k_l} \rightarrow e$ as $l \rightarrow \infty$. Hence $s_t = e$, which is a contradiction.

It follows from Claim that there exists a strictly increasing sequence $\{n_l\}_{l \in \mathbb{N}} \subset \mathbb{N}$ such that $m_{n_i} < m_{n_{i+1}}$ for each $i \in \mathbb{N}$. Therefore, we can set $i_l = n_l$ and $j_l = m_{n_l} + k_{n_l}$ for each $l \in \mathbb{N}$. Then we get the strictly increasing sequences $\{i_l\}_{l \in \mathbb{N}}$ and $\{j_l\}_{l \in \mathbb{N}}$ such that $\lim_l a_{i_l, j_l} = e$. $\square$

It follows from Lemma 5.1, we have the following theorem, which was proved in [18].

Corollary 5.2. *A rectifiable space G is Fréchet-Urysohn if and only if it is strongly Fréchet-Urysohn.*

Lemma 5.3. *Let G be an α_4 -rectifiable space. If G is a sequential space then G is strongly Fréchet-Urysohn.*

Proof. It follows from Corollary 5.2 that it suffices to show that G is Fréchet-Urysohn. Suppose that G is non-Fréchet-Urysohn. Then there exists a subset A of G such that $\hat{A} \setminus \hat{A} \neq \emptyset$, where the set $\hat{A}$ is all the limit points of convergent sequences in A . Take a point $x \in \hat{A} \setminus \hat{A}$. Without loss of generality, we may assume the $x = e$.

Since $e \in \hat{A}$, there exists a sequence $\{x_n\}_{n=1}^\infty \subset \hat{A}$ such that the sequence $\{x_n\}_{n=1}^\infty$ converges to e . For each $n \in \mathbb{N}$, there exists a sequence $\{x_{nj}\}_{j=1}^\infty \subset A$ such that the sequence $\{x_{nj}\}_{j=1}^\infty$ converges to x_n . Since G is a rectifiable space, the sequence $\{q(x_n, x_{nj})\}_{j=1}^\infty$ converges to $q(x_n, x_n) = e$ as $j \rightarrow \infty$. Moreover, since G

is an α_4 -rectifiable space, there are an increasing sequence $\{n_k\}_{k=1}^\infty$ and a sequence $\{j(n_k)\}_{k=1}^\infty$ such that $\{q(x_{n_k}, x_{n_k j(n_k)})\}_{k=1}^\infty$ converges to e . Then we have

$$x_{n_k j(n_k)} = p(x_{n_k}, q(x_{n_k}, x_{n_k j(n_k)})) \rightarrow p(e, e) = e \text{ as } k \rightarrow \infty.$$

However, we have $e \notin \hat{A}$, which is a contradiction. $\square$

It follows from Lemmas 5.1, 5.3 and Corollary 5.2 that we have the following theorem.

Theorem 5.4. *Let G be a sequential rectifiable space. Then the following conditions are equivalent:*

- (1) *The space G is an α_4 -space;*
- (2) *The space G is an AS-space;*
- (3) *The space G is Fréchet-Urysohn;*
- (4) *The space G is strongly Fréchet-Urysohn.*

Corollary 5.5. [15] *If G is a weakly first-countable rectifiable space, then G is first-countable and hence it is metrizable.*

Proof. It is well known that a weakly first-countable space is a sequential α_4 -space, by Lemma 5.3, G is a Fréchet-Urysohn space, then G is first-countable since a Fréchet-Urysohn weakly first-countable space is first-countable. Hence G is metrizable. $\square$

6. METRIZABILITIES OF RECTIFIABLE SPACES

In [18], F.C. Lin and R.X. Shen posed the following question:

Question 6.1. [18] *Is every sequential rectifiable space with a point-countable k -network² a paracompact space?*

In this section, we shall give a partial answer for Question 6.1. Moreover, we also discuss the metrizability of rectifiable spaces.

Let X be a space and $x \in X$. The space X has property $P(x, U)$ [23] if $U \subset X$, $\{x_i : i \in \mathbb{N}\} \subset U$, $x_i \rightarrow x$ as $i \rightarrow \infty$ and $x_i \neq x_j$ if $i \neq j$, then there is $\Gamma = \{x(n, k) : n, k \in \mathbb{N}\} \subset U$ such that $x(n, k) \rightarrow x_n$ as $k \rightarrow \infty$, $t : \mathbb{N}^2 \rightarrow \Gamma$ is a bijection, where $t(n, k) = x(n, k)$ and $\Gamma \cup \{x_i : i \in \mathbb{N}\} \cup \{x\}$ is a closed subset of X homeomorphic to S_2 .

Lemma 6.2. [23] *A sequential non-Fréchet-Urysohn space with a point-countable k -network contains a closed copy of S_2 .*

Lemma 6.3. [18] *Let G be a rectifiable space. Then G contains a (closed) copy of S_ω if and only if G has a (closed) copy of S_2 .*

Lemma 6.4. *Let G be a non-Fréchet-Urysohn sequential rectifiable space with point-countable k -network. Then for any $x \in G$ and any open $U \subset G$, G has the property $P(x, U)$.*

² Let $\mathcal{P}$ be a family of subsets of a space X . The family $\mathcal{P}$ is called a k -network [20] if whenever K is a compact subset of X and $K \subset U \in \tau(X)$, there is a finite subfamily $\mathcal{P}' \subset \mathcal{P}$ such that $K \subset \bigcup \mathcal{P}' \subset U$.

Proof. By Lemma 6.2, there exists a closed subset of G homeomorphic to S_2 . It follows from Lemma 6.3, G contains a closed subset homeomorphic to S_ω . Let $S_\omega = \{y(n, k) : n, k \in \mathbb{N}\} \cup \{e\}$, where $y(n, k) \rightarrow e$ as $k \rightarrow \infty$ and $y(n, k) \neq y(l, m)$ if $(n, k) \neq (l, m)$. We may assume that S_ω is a closed subset of G . Let U be open in G , $\{x_i : i \in \mathbb{N}\} \subset U$, $x_i \rightarrow x$ as $i \rightarrow \infty$ and $x_i \neq x_j$ if $i \neq j$. For any $i, k \in \mathbb{N}$, put $x(i, k) = p(x_i, y(i, k))$. Since $y(i, k) \rightarrow e$ as $k \rightarrow \infty$, it follows that $x(i, k) \rightarrow p(x_i, e) = x_i$ as $k \rightarrow \infty$. For every $i \in \mathbb{N}$, we can choose a $k_i \in \mathbb{N}$ such that $\{x(i, k) : i \in \mathbb{N}, k \geq k_i\} \subset U$ and $x(i, k) \neq x(i', k')$ if $(i, k) \neq (i', k')$ and $k \geq k_i, k' \geq k_{i'}$. Then $\{x_i : i \in \mathbb{N}\} \cup \{x(i, k) : i \in \mathbb{N}, k \geq k_i\} \cup \{x\}$ is a closed subset in G and homeomorphic to S_2 . If not, there exists a sequence $\{x(i_j, k_j)\}_{j=1}^\infty$ converging to some point $b \in G$ such that $i_j \neq i_{j'}$, if $j \neq j'$. Therefore, we have

$$y(i_j, k_j) = q(x_{i_j}, p(x_{i_j}, y(i_j, k_j))) = q(x_{i_j}, x(i_j, k_j)) \rightarrow q(x, b) \text{ as } j \rightarrow \infty.$$

However, the set $\{y(i_j, k_j) : j \in \mathbb{N}\}$ is closed and discrete in G , which is a contradiction. $\square$

Lemma 6.5. [23] *Let X be a sequential space with a point-countable k -network such that for any $x \in X$ and $U \subset X$ the property $P(x, U)$ holds. Then for any $\alpha < \omega_1, x \in X, U \subset X$ open in X the following property holds:*

$Q(\alpha, x, U)$: *If $\{x_i : i \in \mathbb{N}\} \subset U, x_i \rightarrow x$ as $i \rightarrow \infty$ then there is $Q \subset U$ such that $\overline{Q}$ is countable, $\overline{Q} \setminus \{x\} = U, x \in [Q]_\alpha, x \notin [Q]_\beta$ for each $\beta < \alpha$.*

Lemma 6.6. [18] *Let G be a sequential rectifiable space. If G has a point-countable k -network, then G is metrizable if and only if G contains no closed copy of S_2 .*

Theorem 6.7. *Let G be a sequential rectifiable space with a point-countable k -network. If $so(G) < \omega_1$, then G is metrizable.*

Proof. Suppose that $so(G) = \alpha$.

Claim: The space G is Fréchet-Urysohn.

Suppose not, it follows from Lemmas 6.4 and 6.5 that G has property $Q(\alpha + 1, e, G)$. Clearly, since G has the property $Q(\alpha + 1, e, G)$, we have $so(G) \geq \alpha + 1 > \alpha$, which is a contradiction.

It follows from the claim that G is a Fréchet-Urysohn rectifiable space, and hence G contains no closed copy of S_2 . Since G is a Fréchet-Urysohn rectifiable space with a point-countable k -network, the space G is metrizable by Lemma 6.6. $\square$

Proposition 6.8. Let $\mathcal{P}$ be a topological property that is productive and preserved by continuous maps. Then the following are equivalent for a rectifiable space G .

- (i) Every subset with the property $\mathcal{P}$ of G has countable pseudocharacter.
- (ii) Every subset with the property $\mathcal{P}$ of G has regular G_δ -diagonal³.

Proof. (ii) $\rightarrow$ (i) obvious.

(i) $\rightarrow$ (ii). Let A be a subset of G and have the property $\mathcal{P}$. Since $q : G \times G \rightarrow G$ is continuous and the property $\mathcal{P}$ is productive and preserved by continuous maps, then $q(A \times A) = B$ is a subset of G and B has the property $\mathcal{P}$. Then $e \in B$ since $q(x, x) = e$. Therefore, e is a G_δ -set of B , let $\{U_n : n \in \mathbb{N}\}$ be a family of countable open subsets with $e \in U_n$ and $\overline{U_{n+1}} \subset U_n$. Then $\Delta = \{(x, x) : x \in A\} = \bigcap_{n \in \mathbb{N}} q^{-1}(U_n) = \bigcap_{n \in \mathbb{N}} \overline{q^{-1}(U_n)}$. In fact, let $(x, y) \in \bigcap_{n \in \mathbb{N}} q^{-1}(U_n)$. For

³A space X is said to have a *regular G_δ -diagonal* if the diagonal $\Delta = \{(x, x) : x \in X\}$ can be represented as the intersection of the closures of a countable family of open neighborhoods of Δ in $X \times X$.

each $n \in \mathbb{N}$, we have $(x, y) \in q^{-1}(U_n)$, and hence $q(x, y) \in U_n$, which follows that $\pi_2(\varphi(x, y)) \in U_n$, $\pi_2(x, y') \in U_n$, where $\varphi(x, y) = (x, y')$ and $y' \in U_n$ for each $n \in \mathbb{N}$. Therefore, $y' = e$. Since $\varphi(x, x) = e$, $\varphi(x, y) = e$ and φ is one-to-one, we have $x = y$. Therefore $\Delta = \bigcap_{n \in \mathbb{N}} q^{-1}(U_n)$. Since $\overline{q^{-1}(U_{n+1})} \subset q^{-1}(\overline{U_{n+1}}) \subset q^{-1}(U_n)$, then we can see that A has a regular G_δ -diagonal. $\square$

It is well known that a countably compact (compact) space with a G_δ -diagonal is metrizable, we have the following.

Corollary 6.9. *The following are equivalent for a rectifiable space G .*

- (i) *Every compact (countably compact) subset is first-countable.*
- (ii) *Every compact (countably compact) subset is metrizable.*

Corollary 6.10. *Let G be a rectifiable space of countable pseudocharacter. Then G has a regular G_δ -diagonal.*

7. COMPACTIFICATIONS OF RECTIFIABLE SPACES

In this section, we assume that all spaces are Tychonoff.

Note that a rectifiable space is metrizable if its π -character is countable [15], by the same proof of [19, Lemma 2], we can prove the following.

Lemma 7.1. *Let G be a non-locally compact rectifiable space. If for each $y \in Y = bG \setminus G$, there exists an open neighborhood $U(y)$ of y such that every countably compact subset of $U(y)$ is metrizable and $\pi\chi(U(y)) \leq \omega$, then G is metrizable and locally separable.*

A space X is called having the property (*): if the cardinality of X is Ulam non-measure, then X is weakly HN-complete⁴. A paracompact space has the property (*) since a paracompact space with Ulam non-measurable cardinality is HN-complete [7, 13], and hence it is weakly HN-complete.

Proposition 7.2. *Let G be a non-locally compact rectifiable space with property (*). If for each $y \in Y = bG \setminus G$, there exists an open neighborhood $U(y)$ of y such that (i) every compact subset of $U(y)$ is a G_δ -subset of $U(y)$; (ii) every countably compact or Lindelöf p -subspace of $U(y)$ is metrizable. Then G, bG are separable and metrizable.*

Proof. From condition (ii), we can see that Y is not locally countably compact, otherwise G is closed in bG and is compact.

By [6, Theorem 3.1], Y is pseudocompact or Lindelöf.

Case 1. The space Y is pseudocompact. Then Y is first-countable since each singleton of Y is a G_δ -set. Since Y is not locally countably compact, the rectifiable space G is locally separable and metrizable by Lemma 7.1. Y is Lindelöf [16] since G is of countable type. Therefore, Y is compact, and hence G is locally compact, which is a contradiction.

Case 2. The space Y is Lindelöf. Since Y is a space of countable pseudocharacter, it follows that the cardinality of Y is Ulam non-measurable [7]. The space G is

⁴A space X is *weakly HN-complete* if the remainder Z of X in the Čech-Stone compactification βX of X is a space of point-countable type.

not locally compact, then G is nonwhere locally compact since G is homogeneous. It follows that G is a remainder of Y , so the cardinality of G is also Ulam non-measurable [7]. Then G is weakly HN-complete. By [5, Theorem 4], each G_δ -point of Y is a point of bisequentiality of Y , it follows that $\pi\chi(Y) \leq \omega$. Therefore, G is locally separable and metrizable by Lemma 7.1. We write $G = \bigoplus_{\alpha \in A} G_\alpha$, where G_α is a separable metrizable subset for each $\alpha \in A$. Let $\eta = \{G_\alpha : \alpha \in A\}$, and let F be the set of all points of bG at which η is not locally finite. Since η is discrete in X . Then $F \subset bG \setminus G$. It is easy to see that F is compact, we can find finitely many closed neighborhoods that satisfy (ii) to cover F , hence F is separable and metrizable, thus F has a countable network. Put $M = Y \setminus F$. For each point $y \in M$, there is an open neighborhood O_y satisfying (ii) in bG such that $\overline{O_y} \cap F = \emptyset$. Since η is discrete, the set $\overline{O_y}$ meets at most finitely many G_α . Let $L = \bigcup \{G_\alpha : G_\alpha \cap \overline{O_y} \neq \emptyset\}$. Then L is separable metrizable. It follows that $\overline{L} \setminus L$ is a remainder of L , and hence it is a Lindelöf p -space by [2, Theorem 2.1]. $Cl_Y(O_y) \subset \overline{L} \setminus L$, then $Cl_Y(O_y)$ is a Lindelöf p -space, hence it is separable and metrizable and $Y \setminus F$ is locally separable metrizable. Since F is compact, there are finite many $\{U(y_i) : i \leq k\}$ that satisfy (i) and cover F . Moreover, since each compact subset of $U(y_i) (i \leq k)$ is a G_δ -set, the set F is a G_δ -set in $\bigcup \{U(y_i) : i \leq k\}$. We write $F = \bigcap V_n$ with V_n open in Y and $Cl_Y(V_{n+1}) \subset V_n$. Let $K_1 = Y \setminus V_1, K_n = Cl(V_{n-1}) \setminus V_n (n > 1)$. Since Y is Lindelöf and K_n is closed in Y , each K_n is Lindelöf and locally separable metrizable. Therefore, K_n has a countable base for each n . Since $Y = F \cup (\bigcup \{K_n : n \in \mathbb{N}\})$, it follows that Y has a countable network. Then $c(Y) \leq \omega$, hence $c(G) \leq \omega$. Since G is a metrizable space with countable Souslin number, the space G is separable and metrizable. It follows that bG is separable and metrizable since G and Y both have countable networks. $\square$

Recall that the space X has a *quasi- G_δ -diagonal* provided there is a sequence $\{\mathcal{G}(n) : n \in \mathbb{N}\}$ of collections of open sets with property that, given distinct points $x, y \in X$, there is some n with $x \in st(x, \mathcal{G}(n)) \subset X \setminus \{y\}$. Obviously, “ X has a G_δ -diagonal” implies “ X has a quasi- G_δ -diagonal”.

Theorem 7.3. *Let G be a non-locally compact, paracompact rectifiable space, and $Y = bG \setminus G$ have locally quasi- G_δ -diagonal. Then G and bG are separable and metrizable.*

Proof. By [8, Proposition 2.3], for $y \in Y$, there exists an open neighborhood $U(y)$ such that each compact subset of $U(y)$ is a G_δ -set and every countably compact subset of $U(y)$ is metrizable. Moreover, every Lindelöf p -subspace of $U(y)$ is metrizable by [17, Corollary 3.6]. Then G and bG are separable and metrizable by Proposition 7.2 $\square$

Corollary 7.4. [6] *Let G be a non-locally compact, paracompact rectifiable space, and $Y = bG \setminus G$ have a G_δ -diagonal. Then G and bG are separable and metrizable.*

Proposition 7.5. *Let G be a non-locally compact rectifiable space. If for each $y \in Y = bG \setminus G$, there exists an open neighborhood $U(y)$ of y such that (i) $\pi\chi(U(y)) \leq \omega$; (ii) every countably compact or Lindelöf p -subspace of $U(y)$ is metrizable; (iii) every compact subset of $U(y)$ is a G_δ -set of $U(y)$. Then G, bG are separable and metrizable.*

Proof. By Lemma 7.1, G is metrizable and locally separable. Similar to the proof of Proposition 7.2, G and bG are separable and metrizable. $\square$

A space with point-countable base satisfies (i), (ii) [14, Corollary 7.11(ii)] and (iii) in Proposition 7.5.

Corollary 7.6. *Let G be a non-locally compact rectifiable space, and $Y = bG \setminus G$ have locally point-countable base. Then G and bG are separable and metrizable.*

By [8, Proposition 2.1] and [14, Corollary 8.3(ii)], a space with a $\delta\theta$ -base⁵ satisfies (i), (ii) and (iii) in Proposition 7.5.

Corollary 7.7. *Let G be a non-locally compact rectifiable space, and $Y = bG \setminus G$ have locally $\delta\theta$ -base. Then G and bG are separable and metrizable.*

By [14, Corollary 10.7(ii), Theorem 10.6], a γ -space⁶ satisfies (i), (ii) and (iii) in Proposition 7.5.

Corollary 7.8. *Let G be a non-locally compact rectifiable space, and $Y = bG \setminus G$ be a locally γ -space. Then G and bG are separable and metrizable.*

Acknowledgements. We wish to thank the reviewers for the detailed list of corrections, suggestions to the paper, and all her/his efforts in order to improve the paper.

REFERENCES

- [1] A.V. Arhangel'skiĭ, Mappings and spaces, *Russian Math. Surveys*, **21**(1966), 115–162.
- [2] A.V. Arhangel'skiĭ, *Remainders in compactifications and generalized metrizability properties*, Topology Appl., 150(2005), 79–90.
- [3] A.V. Arhangel'skiĭ, *More on remainders close to metrizable spaces*, Topology Appl., 154(2007), 1084–1088.
- [4] A.V. Arhangel'skiĭ, M. Tkachenko, *Topological Groups and Related Structures*, Atlantis Press and World Sci., 2008.
- [5] A.V. Arhangel'skiĭ, G_δ -points in remainders of topological groups and some addition theorems in compacta, Top. Appl., 156(2009), 2013–2018.
- [6] A.V. Arhangel'skiĭ, M.M. Choban, *On remainders of rectifiable spaces*, Topology Appl., **157**(2010), 789–799.
- [7] A.V. Arhangel'skiĭ, V. Ponomarev, *Fundamentals of General Topology in Problems and Exercises*, Reidel. Dordrecht, 1984 (translated from Russia).
- [8] H. Bennett, R. Byerly, D. Lutzer, *Compact G_δ -sets*, Top. Appl., 153(2006), 2169–2181.
- [9] T. Banach, *On topological groups containing a Fréchet-Urysohn fan*, Matematychni Studii, **9**(1998), 149–154.
- [10] T. Banach, L. Zdomskyĭ, *The topological structure of (homogeneous) spaces and groups with countable cs^* -character*, Appl. Gen. Topology, **5**(2004), 25–48.
- [11] M.M. Čoban, *On topological homogeneous algebras*, in: Interim Reports of Prague Topol. Symp. II, Prague, 1987, 25–26.
- [12] M.M. Čoban, *The structure of locally compact algebras*, Serdica, **18**(1992), 129–137.
- [13] R. Engelking, *General Topology* (revised and completed edition), Heldermann Verlag, Berlin, 1989.
- [14] G. Gruenhage, *Generalized metric spaces*, K. Kunen, J.E. Vaughan eds., Handbook of Set-Theoretic Topology, North-Holland, (1984), 423–501.
- [15] A.S. Gul'ko, *Rectifiable spaces*, Topology Appl., **68**(1996), 107–112.

⁵_a $\delta\theta$ -base for a space X is a base $\mathcal{B} = \cup\{\mathcal{B}(n) : n \geq 1\}$ with the additional property that U is open and $x \in U$, then there is some $n = n(x, U)$ with properties that (a) some $B \in \mathcal{B}(n)$ has $x \in B \subset U$, and (b) $\text{ord}(x, \mathcal{B}(n)) \leq \omega$.

⁶A space (X, τ) is a γ -space if there exists a function $g : \omega \times X \rightarrow \tau$ such that (i) $\{g(n, x) : n \in \omega\}$ is a base at x ; (ii) for each $n \in \omega$ and $x \in X$, there exists $m \in \omega$ such that $y \in g(m, x)$ implies $g(m, y) \subset g(n, x)$.

- [16] M. Henriksen, J. Isbell, *Some properties of compactifications*, Duke Math. J. 25(1958), 83-106.
- [17] R. Hodel, *Metrizability of topological spaces*, Pacific J. Math., 55(1974), 441-459.
- [18] F.C. Lin, R.X. Shen, *On rectifiable spaces and paratopological groups*, Topology Appl., to appear.
- [19] C. Liu, *Remainders in compactification of topological groups*, Top. Appl., 156(2009), 849-854.
- [20] P. O'Meara, *On paracompactness in function spaces with the compact-open topology*, Proc. Amer. Math. Soc., **29**(1971), 183-189.
- [21] T. Nogura, D. Shakhmatov, *Amalgamation of convergent sequences in locally compact groups*, C. R. Acad. Sci. Paris, Serie 1, **320**(1995), 1349-1354.
- [22] E. Pentsak, *On manifolds modeled on direct limits of C -universal ANR's*, Matematychni Studii, **5**(1995), 107-116.
- [23] A. Shibakov, *Metrizability of sequential topological groups with point-countable k -networks*, Proc. Amer. Math. Soc., **126**(1998), 943-947.
- [24] V.V. Uspenskij, *The Maltsev operation on countably compact spaces*, Comments. Math. Univ. Carolin., **30**(1989), 395-402.
- [25] V.V. Uspenskij, *Topological groups and Dugundji compacta*, Mat. Sb. **180**(1989), no. 8, 1092-1118 (Russian); English transl. in: Math. USSR-Sb. **67**(1990), no. 2, 555-580.

(FUCAI LIN): DEPARTMENT OF MATHEMATICS AND INFORMATION SCIENCE, ZHANGZHOU NORMAL UNIVERSITY, ZHANGZHOU 363000, P. R. CHINA
E-mail address: `linfucal2008@yahoo.com.cn`

(CHUAN LIU): DEPARTMENT OF MATHEMATICS, OHIO UNIVERSITY ZANESVILLE CAMPUS, ZANESVILLE, OH 43701, USA
E-mail address: `liuc1@ohio.edu`

(SHOU LIN): INSTITUTE OF MATHEMATICS, NINGDE TEACHERS' COLLEGE, NINGDE, FUJIAN 352100, P. R. CHINA
E-mail address: `linshou@public.ndptt.fj.cn`